

News Articles Videos Guest Columns Blog Browse Topics About Us E-mail Alert

INSIDE SCIENCE NEWS SERVICE

Cosmic Experiment Aims To Close Loophole In Quantum Theory

Like Share 5 Tweet 13 **8+1** 3



Image credit: ESO/M. Kornmesser | http://bit.ly/1kyq6q0

Distant guasars could help confirm "spooky action" between particles.

Originally published: Mar 5 2014 - 12:15pm

By: Charles Q. Choi, ISNS Contributor

(ISNS) -- An experiment of cosmic proportions, looking at some of the most distant visible corners in the universe, could help close what may be the last major loophole in quantum physics, or shake it to its very foundations.

In the bizarre realm of quantum physics, two or more particles can get linked so they stay in sync instantaneously no matter how far apart they are. Albert Einstein derisively called this seemingly impossible connection "spooky action at a distance" - scientists nowadays give it the name guantum entanglement. Einstein believed that guantum entanglement could be explained on a deeper level by the more intuitive laws of classical physics.

In the classical picture, two objects in different regions of space could not influence each other faster than the speed of light. Also, every particle would have well-defined properties at every moment in time -- classical mechanics-based, "hidden variables" that would dictate the strangely synchronized entangled behavior between the particles.

Fifty years ago, physicist John Bell devised a mathematical formula that predicted what scenarios would occur if the counterintuitive predictions of quantum physics were governed by these classical hidden variables. Experiments on Bell's theorem have supported a purely quantum physics picture, rejecting the existence of hidden variables and showing particles are linked more strongly than one would expect under the laws of classical physics.

However, researchers have also identified major potential loopholes in Bell's theorem.

Two have been closed, but a third remains, "one known as 'setting independence' or sometimes colorfully called the 'free will' loophole," said theoretical physicist David Kaiser at MIT. "It's a really crazy-sounding loophole, but it turns out it's the easiest way to fake an outcome in tests of Bell's theorem."

In this scenario, the two detectors measuring the entangled particles have a shared history, via an event, information or third party they have in common. This could link them and lead to

FROM THE CURRENTS BLOG

In Space, Could Rotation Make The Immune System Stronger?

In Speed Skating, Other Sports, Air Resistan Plays A Big Role

The British Invasion In Space

2013 Year In Review: Inside Science

Turning Solar Storms Into Music

RELATED ARTICLES

Building Unbreakable Codes Beyond The Bou Of Earth

Proposed Time Machine Could Also Clone Ob

Objects Beyond Cosmic Voids Not As Bright / They Appear

Where Did The Universe's Magnetism Come From?

Mystery Force May Be Due To Mirrors

MORE FROM THIS DEPARTMENT

Building Unbreakable Codes Beyond The Bou Of Earth

Proposed Time Machine Could Also Clone Ob

Mystery Force May Be Due To Mirrors

Jupiter's Melting Heart Sheds Light on Myste Exoplanet

To Date, Particle Supercollider Detects No Evidence Of Dark Matter

EDITOR'S PICKS

Electronic Devices That Dissolve In Your Bod

Black Hole Cores May Not Be Infinitely Dense

Physicist Proposes New Way To Think About Intelligence

The Science Behind 'Beatboxing'

Every Black Hole Contains a New Universe

Advertising

 \square

biased results. Therefore, a scientist testing Bell's theorem would not have complete control in choosing what each detector measures.

To solve this loophole, Kaiser and his colleagues have proposed looking for answers from the most remote corners of the known cosmos.

The experiment they propose relies on the fact that the universe has been expanding ever since the Big Bang happened nearly 14 billion years ago. As such, objects can be far enough away from each other to have been out of contact since the beginning of the cosmos, with no way for any signal to have ever reached from one to the other. This makes it possible to test the last loophole in Bell's theorem for a so-called "Cosmic Bell" experiment.

The experiment would involve a pair of telescopes, aimed at opposite sides of the sky at different quasars -- supermassive black holes up to billions of times the mass of the sun, which release extraordinarily large amounts of light as they devour matter.

The experiment would create two entangled particles here on Earth and send one of the two to both telescopes. The particles would be measured by detectors at the telescope, machines that would rely on information from the two unconnected quasars to determine which properties they would measure of the two entangled particles.

The scientists reason that because each detector's settings are controlled by quasars that have shared no history since the universe was born, it should be impossible for these detectors to be part of a conspiracy to skew their results. If the experiment discovered the measurements of the entangled particles matched each other more than predicted by the laws of classical physics, that should close the "free will" loophole.

Although the researchers expect this experiment to verify quantum theory's predictions, if the test finds otherwise, "that would be a win also," Kaiser said. "It could mean that we have to change quantum theory, the fundamental theory governing matter. Or it could mean that there could be weird activity in the very earliest moments of the Big Bang."

Kaiser said that the experiment is possible using modern technology.

The distance between each telescope and the source of the entangled particles needs to be on the order of 50 kilometers. Scientists have already carried out experiments with entangled particles 144 kilometers apart, between the two Canary Islands of La Palma and Tenerife off the northwest coast of Africa, a feat that was announced in 2007.

"It turns out the Canary Islands have some of the largest optical telescopes in the world, so maybe we could do it there," Kaiser said.

High energy physicist Warren Huelsnitz with the MiniBooNE experiment at Fermilab in Batavia, III., who did not take part in this research, noted the experiment would be very challenging — scientists would have to make sure the light the telescopes measure really came from the quasars, as opposed to light pollution on Earth, or scattered starlight.

However, "If Bell's inequality is not violated in the 'Cosmic Bell' experiment, then that would be truly amazing and it would set quantum mechanics, and perhaps all of physics, on its head," Huelsnitz said. Nevertheless, Huelsnitz predicts that the results will likely be consistent with previous tests of Bell's theorem.

Even closing this loophole "will not entirely rule out hidden-variables theories," cautioned theoretical physicist Sabine Hossenfelder at the Nordic Institute for Theoretical Physics in Stockholm, Sweden, who did not participate in this study. For instance, there are so-called superdeterministic hidden variables that Bell's theorem cannot be used to test. However, she is now investigating whether experiments could show whether those hidden variables exist or not as well.


Kaiser and his colleagues Jason Gallicchio and Andrew Friedman will detail <u>their findings</u> in the journal <u>Physical Review Letters</u>.

Blocked Plug-in

Charles Q. Choi is a freelance science writer based in New York City who has written for The New York Times, Scientific American, Wired, Science, Nature, and many other news outlets. He tweets at <u>@cqchoi</u>.

Science category: Astronomy Physics Space/Cosmos News section: Inside Science News Service		
Tags:	Quantum Physics entanglement hidden variables Einstein Bell's theorem	
0 Comme	ents InsideScience	D Login -
Sort by Newest *		Favorite ★
	Start the discussion	

Be the first to comment.



Reprint rights | Commenting policy

About Inside Science | Contact us | Privacy policy | Reprint rights | ©2014 American Institute of Physics